

African Penguin *Spheniscus demersus* foraging on juvenile fish in jellyfish tentacles

Jacqueline S. Glencross^{1*}, Amí Jewell², Jennifer L. Grigg², Alistair McInnes³,
Andrew S. Brierley^{1†} & Richard B. Sherley^{2,4}

* Correspondence author. Email: jg287@st-andrews.ac.uk

¹ Scottish Oceans Institute, University of St Andrews, Fife, KY16 8LB, UK;

² Environment and Sustainability Institute, University of Exeter, Cornwall,
TR10 9FE, UK;

³ BirdLife South Africa, Cape Town 8001, South Africa;

⁴ Department of Biodiversity and Conservation Biology, University of the Western
Cape, Bellville, 7535, South Africa.

† Deceased.

JS Glencross ORCID: 0000-0001-8581-5417; JL Grigg ORCID: 0000-0002-0576-9359;
A McInnes ORCID: 0000-0002-9125-9629; AS Brierley ORCID: 0000-0002-6438-6892;
RB Sherley ORCID: 0000-0001-7367-9315.

Abstract

Technological advances have enabled the observation of foraging behaviour in wild marine animals. We can observe where they go, how deep they dive, how much energy they expend, and with the use of animal-borne cameras we can capture specific foraging behaviours. Here we describe a newly observed foraging behaviour in African Penguins *Spheniscus demersus* in which they target fish located in jellyfish tentacles. As animal-borne cameras have only been deployed on African Penguins since 2015 it is unclear whether this behaviour is novel or previously unobserved. The behaviour appears to be opportunistic and beneficial to the African Penguins as it enables them to catch stationary prey. As the availability of the African Penguin's traditional schooling fish prey changes in the southern Benguela Current ecosystem due to climate change and overfishing, opportunistic foraging strategies like this could enable African Penguins to supplement foraging success at a low energetic cost.

Introduction

African Penguins *Spheniscus demersus* are endemic to southern Africa where they breed in South Africa and Namibia. Their numbers have declined by >80% since the first comprehensive census in the 1970s (Sherley *et al.* 2020). As a consequence, the species was listed as 'Endangered' on the IUCN Red List in 2010 (Sherley *et al.* 2020) and may soon meet the criteria for being listed as 'Critically Endangered' (Sherley *et al.* 2024). The common prey species of the African Penguin, determined through stomach flushing on breeding adults, have traditionally been Southern African Anchovy *Engraulis capensis* (hereafter 'Anchovy') and Sardine *Sardinops sagax* (Crawford *et al.* 2011).

However, following the collapse of Sardine stocks in the northern Benguela in the 1960s and 1970s, the Namibian population of African Penguins switched to feeding almost exclusively on Bearded Goby *Sufflogobius bibarbatus* (Ludynia *et al.* 2010). In the southern Benguela Current, the Sardine stock has been in a poor state for the last c. 15 years following a combination of climate change and fishing pressure (Coetzee *et al.* 2008; Coetzee *et al.* 2022), and although Anchovy remains abundant, its availability to seabirds may have changed following an eastward displacement of spawning habitat (Crawford *et al.* 2019; Mhlongo *et al.* 2015).

The change in forage fish distributions and abundances in the Benguela ecosystem, alongside declining African Penguin populations, has led to substantial research interest in how the availability of prey resources to African Penguins may be changing (e.g. Crawford *et al.* 2006, 2011; Campbell *et al.* 2019). The colony on Robben Island, South Africa, is the focus of one of the longest running monitoring projects on African Penguins (Crawford *et al.* 2006; Leith *et al.* 2022). In addition to monitoring metrics like breeding success and chick condition on land (e.g. Campbell *et al.* 2019; Sherley *et al.* 2013), we have deployed GPS dive loggers on breeding African Penguins since 2008, providing important information for management, especially where they are foraging and how this varies through time (e.g. Campbell *et al.* 2019). However, at this time, dive data allow only for inference on when African Penguins are foraging, but not details about what they are eating.

Many diet analysis techniques for use on seabirds are necessarily highly invasive, e.g. stomach flushing (Wilson 1984). It can be difficult to justify these invasive techniques from a contemporary viewpoint, particularly when working with endangered species like the African Penguin. Fortunately, less invasive methods are increasingly available. The use of camera technology to study seabird diet is becoming widespread (e.g. Gaglio *et al.* 2018b; Mattern *et al.* 2018), with animal-borne video cameras providing valuable information on prey species, when the resolution of footage is high enough (McInnes *et al.* 2017; Watanabe & Takahasi, 2013; Ponganis *et al.* 2000). Furthermore, video cameras have the added benefit of documenting foraging behaviour and success (e.g. Sutton *et al.* 2020; Sutton *et al.* 2021; Watanabe & Takahasi 2013).

This paper reports observations from camera footage collected on African Penguins from Robben Island foraging at sea across four years: 2018, 2019, 2022 and 2023. In particular, we report a behaviour that we do not believe to have been reported in this species before, where individual African Penguins target fish located amongst jellyfish tentacles.

Methods

Adult African Penguins in the guard stage of breeding (with chicks between c. 6–15 days old; Seddon & van Heezik, 1993) were equipped with animal-borne video cameras (2018–19: Replay XD 1080 Mini, total dimensions: 94 x 28 x 23 mm, 65 g, resolution 1920 x 1080 px; 2022–23: TechnoSmArt, Rome, Italy, total dimensions: 70 x 23 x max. 24 mm, 32 g, resolution 1280 x 690 px) for one foraging trip from

Robben Island (33°48'S, 18°22'E) during 2018, 2019, 2022 and 2023. Birds were captured by hand at their nest. On Robben Island there are very few predators, and the African Penguin nests are predominantly under vegetation. Chicks were therefore protected while the adult was being handled. Cameras and time-depth-accelerometer loggers (TechnoSmArt, Rome, Italy; L x W x H = 15 x 9 x 2 mm, mass = 0.7 g) were then attached to the midline of each bird's mid- and caudal portion (respectively) on the back using overlapping layers of waterproof tape (Tesa tape #4651, Beiersdorf AG, Hamburg, Germany; Wilson *et al.* 1997). The tape was then secured with cyanoacrylate adhesive (Loctite 401, USA) to prevent the bird from preening the tape off. Combined mass of the loggers did not exceed 4% of the total mass of the African Penguin. Handling time from capturing the bird to release after attaching devices was maximum 17 minutes (mean: 10.5 minutes, $n = 21$). Cameras were deployed for one foraging trip (usually c. 13 hours), and nests were checked daily to ensure the African Penguins went to sea and that the other adult of the pair had resumed brooding. The birds went to sea for one day and the cameras were retrieved the following morning at the nest. If the bird had not gone to sea on the second day after deployment, the device was removed. On retrieval, cameras and any other devices were removed, then head length, bill length, bill depth (mm) and bird mass (g) were measured and used to gauge the sex of the bird (Campbell *et al.* 2016).

To ensure the footage captured foraging behaviour and to maximise useful battery life, the cameras were set up to turn on approximately one hour after sunrise (when the African Penguins are typically already at sea; Ryan *et al.* 2007). The 2018 and 2019 cameras were limited in their battery life and so were set up to run for 20 minutes, then switch off for an hour before switching back on. This resulted in around one hour of footage. The cameras used in 2022 and 2023 had better battery life so would run continuously (with c. five second gaps in between c. ten minute videos) until the battery died which resulted in around six hours of footage per trip. We labelled and analysed the footage using the software BORIS (an event logging software for video data; Friard *et al.* 2016). For the purpose of this study, prey encounters and prey capture attempts were recorded as point events (time-stamped to the moment that the prey became visible in the footage, or the moment the individual attempted to catch prey, respectively). Prey capture events were defined as successful, unsuccessful, or unknown. Prey encounters were categorised as 'individual fish' (with no shoal in sight), 'shoal', 'squid', or 'other'. It was not possible to measure the exact size of prey, because camera position varied slightly between individuals, but fish seen in videos recorded by one individual could be categorised comparatively as 'large' or 'small'.



Results

Over the four years, we collected foraging footage from 15 African Penguins (a further nine deployments resulted in no video data due to African Penguins not initiating a foraging trip or device failure). The most common prey type was fish (Figure 1a, b, e), typically small forage fish. Of known prey encounters ($n = 280$), 258 were fish with 73.6% of these fish occurring as an individual fish rather than

part of a visible shoal. Other notable prey items include a squid (Figure 1d) and a crustacean (Figure 1c). These are probably Cape Hope (or Chokka) Squid *Loligo reynaudii* and Cape Mantis Shrimp *Pterygosquilla capensis* respectively, as these species have previously been recorded in the diet of penguins and other seabirds off South Africa (e.g. Sherley *et al.* 2013, Gaglio *et al.* 2018a, Connan *et al.* 2016). We also recorded one African Penguin interacting with a small piece of orange rope (Figure 1f). Free-swimming gelatinous zooplankton (hereafter 'jellyfish') were often encountered but were never seen being targeted directly by the African Penguins. The number of jellyfish encountered differed between deployments with many videos not containing a single jellyfish during the period cameras were recording.

Two African Penguins were observed targeting jellyfish that had small fish either entangled or taking refuge within their tentacles. This behaviour typically started with the African Penguin approaching the jellyfish from their underside and plucking the fish out from the tentacles (Figure 2 and 3). In most cases low image quality, and sometimes water visibility, prevented the identification of the gelatinous zooplankton species with which fish were associated. In a few cases we could identify to genus or family level, such as Hydrozoa jellyfish (e.g. *Aequorea* spp.) and other zooplankton taxa including ctenophores or salps. In a few instances, the fish appeared to be surrounded by gelatinous material which we believe to be either a ctenophore or a salp (Figure 2d). This behaviour, targeting fish associated with jellyfish, or in some cases other gelatinous zooplankton, was seen in one individual in 2019 (from $n = 2$ individuals tracked) and one in 2023 ($n = 6$ individuals). Foraging on fish in association with jellyfish was not observed in 2018 ($n = 3$ individuals) or 2022 ($n = 4$ individuals). The 2019 African Penguin was observed inspecting jellyfish 11 times, and on three of those occasions we were able to confirm that the jellyfish had small fish within its tentacles. In 2023, the individual was observed encountering jellyfish 60 times and on 16 of those we could confirm fish within the jellyfish tentacles (Table 1). In most cases, jellyfish only contained one individual small fish, typically forage fish like Anchovy, or pipefish *Syngnathus* sp. (Figure 2c and d, and Figure 3), in which case the African Penguin would catch the fish then continue its dive (duration spent interacting with a single jellyfish = c. 1–2 seconds), typically continuing search behaviour or continue their ascent to the surface. Interactions with jellyfish seemed to occur while ascending from search dives rather than the African Penguins diving to target them specifically. Often jellyfish would be clustered meaning the African Penguin could interact with more than one jellyfish within a dive. Less commonly, the African Penguin encountered a large jellyfish (Figure 2a and b) with multiple fish inside. During these encounters the African Penguin spent up to c. 10 seconds taking the fish out of the tentacles.

Figure 1 (opposite). Images of prey items encountered by African Penguins *Spheniscus demersus*. **a)** Individual forage fish. **b)** School of forage fish. **c)** Crustacean, likely Cape Mantis Shrimp *Pterygosquilla capensis*. **d)** Squid, *Loligo* sp. **e)** Unidentified large fish. **f)** Rope fragment.







Figure 2 (opposite top). African Penguins *Spheniscus demersus* foraging on jellyfish-fish aggregations. **a)** African Penguin approaching a jellyfish with many small fish amongst tentacles. **b)** African Penguin closer to jellyfish with fish (circled) amongst tentacles. **c)** African Penguin catching a pipefish *Syngnathus* sp. from the tentacles of a jellyfish. **d)** African Penguin catching fish in the body of a gelatinous zooplankton.

Figure 3 a–c (opposite bottom). Image sequence of prey capture involving a jellyfish *Aequorea* sp. and fish (pipefish, *Syngnathus* sp.) aggregation. See S1 for full video.

Table 1. Summary of gelatinous zooplankton (jellyfish) encountered on a single foraging trip by two African Penguins *Spheniscus demersus*. *No morphological measurements were taken to identify sex in this individual.

Individual ID	Year	Sex	No. of jellyfish encountered but not interacted with	No. of jellyfish interacted with	No. of prey capture events involving a jellyfish
25_2019R	2019	Unsexed*	2	11	33
30_2023R	2023	Female	324	60	51

Discussion

Here we report a newly described foraging behaviour in the African Penguin whereby they target small fish associating with jellyfish. This behaviour was observed in two different individual African Penguins (out of 15 on which video cameras were deployed) during different years, which implies that it could be used by multiple individuals. The small fish are likely juvenile forage fish, such as Anchovy, due to the timing of the camera footage corresponding with southward juvenile fish movements (Hutchings *et al.* 2002). Juvenile Anchovy or 'recruits', have been commonly identified as the main prey source of the African Penguin in this region (Campbell *et al.* 2019; Crawford *et al.* 2011). Fish within the tentacles could either be jellyfish prey or associated as commensals. Either way, the jellyfish appear to act as natural fish aggregating devices. Large jellyfish are therefore particularly beneficial for African Penguins to investigate as they may contain several, easier to catch prey which require very little energetic expense since there is limited amount of chasing. African Penguins appeared to travel at slower swimming speeds when capturing prey associated with jellyfish compared to normal prey capture attempts and so we hypothesise they will expend less energy (Wilson *et al.* 2010). African Penguins typically interacted with jellyfish while on their ascent from a search dive – meaning the main energetic cost of diving has already been incurred (Wilson *et al.* 2010). This upward angle of approach is consistent with other studies, which showed that Little Penguins *Eudyptula minor* and Brünnich's Guillemot *Uria lomvia* approached jellyfish from the underside when ascending from deeper dives (Sutton *et al.* 2015, Sato *et al.* 2015). Jellyfish were once considered energetic 'dead ends' (Arai 2005; Sommer *et al.* 2002) but have since been documented as crucial prey species for some marine megafauna such as Leatherback Turtles *Dermochelys coriacea* (Houghton *et al.* 2006), as well as supplementary prey for some penguin species (Thiebot *et al.* 2017; Sutton *et al.* 2015), and are increasingly understood to facilitate prey capture in others (Sato *et al.* 2015). Here, we highlight for the first time that foraging on fish in association with jellyfish has been witnessed in the

African Penguin; however, it has previously been documented in other penguins, suggesting that multiple species take advantage of jellyfish-fish aggregations. Indeed, jellyfish-fish aggregations have been targeted by Yellow-eyed Penguin *Megadyptes antipodes* (Mattern 2020), Humbolt Penguin *Spheniscus humboldti* (U. Ellenberg pers. comm.), Brünnich's Guillemot (Sato *et al.* 2015), and in some fish species (Bonaldo *et al.* 2004; Masuda *et al.* 2008).

The fish observed in these jellyfish-fish aggregations tended to be small, typically smaller than the average size of fish being caught by the African Penguins (as seen in the camera footage), either individually or as part of a shoal (Figure 2 compared to Figure 1a). There have been many examples of juvenile fish aggregating within the tentacles of jellyfish as a protection strategy (e.g. Griffin *et al.* 2019; Richardson *et al.* 2009) – albeit unsuccessfully in this African Penguin example. Juvenile fish tend to be of lower energetic content than adults because fish weight is related to fish length by a non-linear power function (Froese 2006). In the European Anchovy *Engraulis encrasicolus*, a close relative of the Southern African Anchovy which is the main prey for African Penguins, energy density is proportional to body length and body mass (Dubreuil & Petitgas 2009). Therefore, we can hypothesise that the fish caught by the African Penguins from the tentacles of jellyfish are of low energetic content.

Foraging on fish in association with jellyfish does not only seem to be linked with low prey availability; in the footage of the individual in 2023, many schools and individual fish were recorded and indeed successfully preyed upon. Rather, this appears to be an opportunistic foraging behaviour which has been used, at least, by a few individuals. We also observed pipefish being fed on (Figure 3), which typically only occur in African Penguin diet when prey availability is low, suggesting they are also low energy prey (Sherley *et al.* 2013). However, pipefish were only confirmed to be predated upon in 2023 (when we also observed forage fish schools) when they were associated with jellyfish. Greater Crested Terns *Thalasseus bergii* breeding at Robben Island have also been observed predated on pipefish (Gaglio *et al.* 2018a). While Greater Crested Terns dive less than 1 m under the surface (Gaglio *et al.* 2018a), pipefish are predominantly found deeper than 1 m in the water column, and so it is possible the terns are also taking advantage of pipefish using jellyfish for refuge.

Both fishes (Anchovy and pipefish) identified associating with jellyfish are known African Penguin prey species (Ludynia *et al.* 2010). The majority of other individual prey species (not associated with jellyfish) identified from the animal-borne camera footage generally matches with the results of diet analysis studies, with forage fish species being the most common prey (Connan *et al.* 2016; Ludynia *et al.* 2010).

While some studies on *Spheniscus* penguins have found cnidaria jellyfish (including *Chrysaora* and *Aequorea* sp.) in their diets (e.g. Thiebot *et al.* 2017), we did not directly observe foraging on jellyfish, only the targeted foraging on fish within their tentacles. In most cases where only jellyfish were seen, there was no

attempt to investigate by the African Penguin, but in cases where fish were in association with the jellyfish, African Penguins swam directly to the jellyfish. For example, the African Penguin in 2023 encountered 324 jellyfish which they did not approach (Table 1). No fish were seen in association with these unapproached jellyfish.

We also documented one observation of an African Penguin investigating a piece of rope, highlighting the potential for ingestion of plastic materials. There has been evidence of plastic ingestion in multiple penguin species including the African Penguin (Brandao *et al.* 2011 Vanstreels *et al.* 2019; Caruso *et al.* 2022). While it was not possible to see from the footage whether the African Penguin in question ingested this piece of rope, it showed that the bird caught the rope in its bill in the same way it would have with a fish.

Camera footage enables detailed diet information to be obtained without using invasive methods, while concurrently collecting data on foraging success, conspecific interactions, and specific foraging behaviours (e.g. McInnes *et al.* 2017; Sutton *et al.* 2020). In this study, animal-borne video camera technology has enabled us to document a newly described opportunistic foraging behaviour in the African Penguin whereby individuals are taking advantage of stationary prey that was either captured or taking refuge within jellyfish tentacles. Although the fish found within the jellyfish tentacles seemed to be small, and potentially have a lower energetic value compared to other available prey, the energy expenditure in catching them is likely low due to the African Penguins not having to chase them. Foraging on fish associating with jellyfish does not seem to only occur as a backup strategy during poor conditions, as plenty of shoals were observed in the footage in 2023, but we recognise that it could become more common during years of low Anchovy (preferred prey source) biomass. We hypothesise that these prey captures are only supplementary to other typical African Penguin foraging behaviour.

Acknowledgments

We thank Robben Island Museum (RIM) for permission to carry out research on the island, accommodation, and ferry transport; in particular, Sabelo Madlala and Inga Sipuka supported the fieldwork. Sue Kuyper and our institutions provided logistical support, and we thank Andrew de Blocq, Taron Morris, Andile Mdluli, Nicholas Ngcathu, and Kiah Tasman for their assistance in the field. This study was conducted under permit from the Department of Forestry, Fisheries and the Environment (DFFE permit numbers: RES2018/18, RES2019/50, RES2022/23 and RES2023-20), in partnership with Robben Island Museum, and under animal ethical clearance from the University of Exeter (Reference numbers: 2018/2404, eCORN001760 and 528178). Funding was provided by the Earthwatch Institute (<http://earthwatch.org/>), the Pew Charitable Trusts (through a Pew Fellows Program in Marine Conservation award to RBS), and the San Diego Zoo Wildlife Alliance. The views expressed in this paper are those of the authors and do not necessarily reflect the views of The Pew Charitable Trusts or any other funder.

References

- Arai, M. N. 2005.** Predation on pelagic coelenterates: a review. *Journal of the Marine Biological Association of the United Kingdom* 85: 523–536. <https://doi.org/10.1017/S0025315405011458>
- Bonaldo, R. M., Krajewski, J. P. & Sazima, I. 2004.** Does the association of young fishes with jellyfishes protect from predation? A report on a failure case due to damage to the jellyfish. *Neotropical Ichthyology* 2: 103–105. <https://doi.org/10.1590/S1679-62252004000200008>
- Brandao, M. L., Braga, K. M. & Luque, J. L. 2011.** Marine debris ingestion by Magellanic penguins, *Spheniscus magellanicus* (Aves: Sphenisciformes), from the Brazilian coastal zone. *Marine Pollution Bulletin* 62: 2246–2249. <https://doi.org/10.1016/j.marpolbul.2011.07.016>
- Campbell, K. J., Farah, D., Collins, S. & Parsons, N. J. 2016.** Sex determination of African penguins *Spheniscus demersus* using bill measurements: method comparisons and implications for use. *Ostrich* 87: 47–55. <https://doi.org/10.2989/00306525.2015.1108371>
- Campbell, K. J., Steinfurth, A., Underhill, L. G., Coetzee, J. C., Dyer, B. M., Ludynia, K., Makhado, A. B., Merkle, D., Rademan, J., Upfold, L. & Sherley, R. B. 2019.** Local forage fish abundance influences foraging effort and offspring condition in an endangered marine predator. *Journal of Applied Ecology* 56: 1751–1760. <https://doi.org/10.1111/1365-2664.13409>
- Caruso, G., Bergami, E., Singh, N. & Corsi, I. 2022.** Plastic occurrence, sources, and impacts in Antarctic environment and biota. *Water Biology and Security* 1: 100034. <https://doi.org/10.1016/j.watbs.2022.100034>
- Coetzee, J. C., De Moor, C. L., van der Lingen, C. D. & Butterworth, D. S. 2022.** A summary of the South African sardine (and anchovy) fishery. MARAM/IWS/2022/Sardine/BG1.
- Coetzee, J. C., Merkle, D., De Moor, C. L., Twatwa, N. M., Barange, M. & Butterworth, D. S. 2008.** Refined estimates of South African pelagic fish biomass from hydro-acoustic surveys: quantifying the effects of target strength, signal attenuation and receiver saturation. *African Journal of Marine Science* 30: 205–217. <https://doi.org/10.2989/AJMS.2008.30.2.1.551>
- Connan, M., Hofmeyr, G. J. & Pistorius, P. A. 2016.** Reappraisal of the trophic ecology of one of the world's most threatened Spheniscids, the African Penguin. *PLOS ONE*. 11: e0159402. <https://doi.org/10.1371/journal.pone.0159402>
- Crawford, R. J. M., Altwegg, R., Barham, B. J., Barham, P. J., Durant, J. M., Dyer, B. M., Geldenhuys, D., Makhado, A. B., Pichegru, L., Ryan, P. G., Underhill, L. G., Upfold, L., Visagie, J., Waller, L. J. & Whittington, P. A. 2011.** Collapse of South Africa's penguins in the early 21st century. *African Journal of Marine Science* 33: 139–156. <https://doi.org/10.2989/1814232X.2011.572377>
- Crawford, R. J. M., Barham, P. J., Underhill, L. G., Shannon, L. J., Coetzee, J. C., Dyer, B. M., Leshoro, T. M. & Upfold, L. 2006.** The influence of food availability on breeding success of African penguins *Spheniscus demersus* at Robben Island, South Africa. *Biological Conservation* 132: 119–125. <https://doi.org/10.1016/j.biocon.2006.03.019>
- Dubreuil, J. & Petitgas, P. 2009.** Energy density of anchovy *Engraulis encrasicolus* in the Bay of Biscay. *Journal of Fish Biology* 74: 521–34. <https://doi.org/10.1111/j.1095-8649.2008.02143.x>
- Friard, O., Gamba, M. & Fitzjohn, R. 2016.** BORIS: a free, versatile open-source event-logging software for video/audio coding and live observations. *Methods in Ecology and Evolution* 7: 1325–1330. <https://doi.org/10.1111/2041-210X.12584>
- Gaglio, D., Cook, T. R., McInnes, A., Sherley, R. B. & Ryan, P. G. 2018a.** Foraging plasticity in seabirds: A non-invasive study of the diet of greater crested terns breeding in the Benguela region. *PLOS ONE* 13: e0190444. <https://doi.org/10.1371/journal.pone.0190444>
- Gaglio, D., Sherley, R. B., Ryan, P. G. & Cook, T. R. 2018b.** A non-invasive approach to estimate the energetic requirements of an increasing seabird population in a perturbed marine ecosystem. *Scientific Reports* 8: 8343. <https://doi.org/10.1038/s41598-018-26647-3>

- Griffin, D. C., Harrod, C., Houghton, J. D. R. & Capellini, I. 2019. Unravelling the macro-evolutionary ecology of fish-jellyfish associations: life in the 'gingerbread house'. *Proceedings of the Royal Society B* 286: 20182325. <https://doi.org/10.1098/rspb.2018.2325>
- Houghton, J. D. R., Doyle, T. K., Wilson, M. W., Davenport, J. & Hays, G. C. 2006. Jellyfish aggregations and Leatherback Turtle foraging patterns in a temperate coastal environment. *Ecology* 87: 1967–1972. [https://doi.org/10.1890/0012-9658\(2006\)87\[1967:JAALTF\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[1967:JAALTF]2.0.CO;2)
- Hutchings, L., Beckley, L. E., Griffiths, M. H., Roberts, M. J., Sunby, S. & van der Lingen, C. 2002. Spawning on the edge: spawning grounds and nursery areas around the southern African coastline. *Marine and Freshwater Research* 53: 307–318. <https://doi.org/10.1071/MF01147>
- Leith, F. W., Grigg, J. L., Barham, B. J., Barham, P. J., Ludynia, K., McGeorge, C., Mdluli, A., Parsons, N. J., Waller, L. J. & Sherley, R. B. 2022. Intercolony variation in reproductive skipping in the African penguin. *Ecology and Evolution* 12: e9255. <https://doi.org/10.1002/ece3.9255>
- Ludynia, K., Roux, J. P., Jones, R., Kemper, J. & Underhill, L. G. 2010. Surviving off junk: low-energy prey dominates the diet of African penguins *Spheniscus demersus* at Mercury Island, Namibia, between 1996 and 2009. *African Journal of Marine Science* 32: 563–572. <https://doi.org/10.2989/1814232X.2010.538151>
- Masuda, R., Yamashita, Y. & Matsuyama, M. 2008. Jack mackerel *Trachurus japonicus* juveniles use jellyfish for predator avoidance and as a prey collector. *Fisheries Science* 74: 276–284. <https://doi.org/10.1111/j.1444-2906.2008.01522.x>
- Mattern, T. 2020. *Modelling marine habitat utilisation by yellow-eyed penguins along their mainland distribution: baseline information*. New Zealand Aquatic Environment and Biodiversity Report No. 243. Fisheries New Zealand.
- Mattern, T., McPherson, M. D., Ellenberg, U., van Heezik, Y. & Seddon, P. J. 2018. High definition video loggers provide new insights into behaviour, physiology, and the oceanic habitat of a marine predator, the yellow-eyed penguin. *Peer J* 6: e5459. <https://doi.org/10.7717/peerj.5459>
- McInnes, A. M., McGeorge, C., Ginsberg, S., Pichegru, L. & Pistorius, P. A. 2017. Group foraging increases foraging efficiency in a piscivorous diver, the African penguin. *Royal Society Open Science* 4: 170918. <https://doi.org/10.1098/rsos.170918>
- Mhlongo, N., Yemane, D., Hendricks, M. & van der Lingen, C. D. 2015. Have the spawning habitat preferences of anchovy (*Engraulis encrasicolus*) and sardine (*Sardinops sagax*) in the southern Benguela changed in recent years? *Fisheries Oceanography* 24: 1–14. <https://doi.org/10.1111/fog.12061>
- Ponganis, P. J., Van Dam, R. P., Marshall, G., Knowner, T. & Levenson, D. H. 2000. Sub-ice foraging behavior of Emperor penguins. *Journal of Experimental Biology* 203: 3275–3278. <https://doi.org/10.1242/jeb.203.21.3275>
- Richardson, A. J., Bakun, A., Hays, G. C. & Gibbons, M. J. 2009. The jellyfish joyride: causes, consequences and management responses to a more gelatinous future. *Trends in Ecology and Evolution* 24: 312–22. <https://doi.org/10.1016/j.tree.2009.01.010>
- Ryan, P. G., Peterson, S. L., Simeone, A. & Grémillet, D. 2007. Diving behaviour of African penguins: do they differ from other *Spheniscus* penguins? *African Journal of Marine Science* 29: 153–160. <https://doi.org/10.2989/AJMS.2007.29.2.1.184>
- Sato, N. N., Kokubun, N., Yamamoto, T., Watanuki, Y., Kitaysky, A. S. & Takahashi, A. 2015. The jellyfish buffet: jellyfish enhance seabird foraging opportunities by concentrating prey. *Biology Letters* 11: 20150358. <https://doi.org/10.1098/rsbl.2015.0358>
- Seddon, P. J. & van Heezik, Y. 1993. Behaviour of the jackass penguin chick. *Ostrich* 64: 8–12. <https://doi.org/10.1080/00306525.1993.9634188>

- Sherley, R. B., Crawford, R. J. M., Blocq, A. D., Dyer, B. M., Geldenhuys, D., Hagen, C., Kemper, J., Makhado, A. B., Pichegru, L., Tom, D., Upfold, L., Visagie, J., Waller, L. J. & Winker, H. 2020. The conservation status and population decline of the African penguin deconstructed in space and time. *Ecology and Evolution* 10: 8506–8516. <https://doi.org/10.1002/ece3.6554>
- Sherley, R. B., Underhill, L. G., Barham, B. J., Barham, P. J., Coetzee, J. C., Crawford, R. J. M., Dyer, B. M., Leshoro, T. M. & Upfold, L. 2013. Influence of local and regional prey availability on breeding performance of African penguins *Spheniscus demersus*. *Marine Ecology Progress Series* 473: 291–301. <https://doi.org/10.3354/meps10070>
- Sommer, U., Stibor, H., Katechakis, A., Sommer, F. & Hansen, T. 2002. Pelagic food web configurations at different levels of nutrient richness and their implications for the ratio fish production:primary production. *Hydrobiologia* 484: 11–20. https://doi.org/10.1007/978-94-017-3190-4_2
- Sutton, G., Pichegru, L., Botha, J. A., Kouzani, A. Z., Adams, S., Bost, C. A. & Arnould, J. P. Y. 2020. Multi-predator assemblages, dive type, bathymetry and sex influence foraging success and efficiency in African penguins. *Peer J* 8: e9380. <https://doi.org/10.7717/peerj.9380>
- Sutton, G. J., Bost, C. A., Kouzani, A. Z., Adams, S. D., Mitchell, K. & Arnould, J. P. Y. 2021. Fine-scale foraging effort and efficiency of Macaroni penguins is influenced by prey type, patch density and temporal dynamics. *Marine Biology* 168: 1–16. <https://doi.org/10.1007/s00227-020-03811-w>
- Sutton, G. J., Hoskins, A. J. & Arnould, J. P. 2015. Benefits of group foraging depend on prey type in a small marine predator, the Little Penguin. *PLOS ONE* 10: e0144297. <https://doi.org/10.1371/journal.pone.0144297>
- Thiebot, J.-B., Arnould, J. P. Y., Gómez-Laich, A., Ito, K., Kato, A., Mattern, T., Mitamura, H., Noda, T., Poupart, T., Quintana, F., Raclot, T., Ropert-Coudert, Y., Sala, J. E., Seddon, P. J., Sutton, G. J., Yoda, K. & Takahashi, A. 2017. Jellyfish and other gelata as food for four penguin species – insights from predator-borne videos. *Frontiers in Ecology and the Environment* 15: 437–441. <https://doi.org/10.1002/fee.1529>
- Vanstreels, R. E. T., Pichegru, L., Pfaff, M. C., Snyman, A., Dyer, B. M., Parsons, N. J., Roberts, D. G., Ludynia, K., Makhado, A. & Pistorius, P. A. 2019. Seashell and debris ingestion by African penguins. *Emu – Austral Ornithology* 120: 90–96. <https://doi.org/10.1080/01584197.2019.1659105>
- Watanabe, Y. Y. & Takahasi, A. 2013. Linking animal-borne video to accelerometers reveals prey capture variability. *Proceedings of the National Academy of Sciences* 110: 2199–2204. <https://doi.org/10.1073/pnas.1216244110>
- Wilson, R. P. 1984. An improved stomach pump for penguins and other seabirds. *Journal of Field Ornithology* 55: 109–112.
- Wilson, R. P., Pütz, K., Peters, G., Culik, B., Scolaro, J. A., Charrassin, J.-B. & Ropert-Coudert, Y. 1997. Long-term attachment of transmitting and recording devices to penguins and other. *Wildlife Society Bulletin* 25: 101–106.
- Wilson, R. P., Shephard, E. L. C., Gómez Laich, A., Frere E., Quintana, F. 2010. Pedalling downhill and freewheeling up; a penguin perspective on foraging. *Aquatic Biology* 8: 193–202. <https://doi.org/10.3354/ab00230>



African Penguin *Spheniscus demersus*,
May 2024. © Jacqui Glencross